

No part of the candidate's evidence in this exemplar material may be presented in an external assessment for the purpose of gaining an NZQA qualification or award.

SUPERVISOR'S USE ONLY

S

93101A



Draw a cross through the box (☒) if you have NOT written in this booklet

+

TOP SCHOLAR



Mana Tohu Mātauranga o Aotearoa
New Zealand Qualifications Authority

Scholarship 2023 Biology

Time allowed: Three hours
Total score: 24

ANSWER BOOKLET

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

Write your answers in this booklet.

Start your answer to each question on a new page. Carefully number each question.

Check that this booklet has pages 2–27 in the correct order. Pages 2–4 are blank and are to be used for planning. Pages 5–27 are lined pages for writing your answers.


Do not write in any cross-hatched area (☒). This area may be cut off when the booklet is marked.

YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.


PLANNING

- inbreeding depression
- loss of habitat
 - territory
- loss of prey species
- & selected species
- Traps, hunting car strikes
- Bacteria

PLANNING



PLANNING



Several different factors have lead to the decline of the lynx population, mostly related to increased human activity in their habitat.

The main factor causing the initial decline of the population was ~~hunting~~ hunting by humans, either trophy hunting or pest control. Even though deliberate hunting was outlawed in the 1970s, indirect killing of lynx's occur through traps and snares by human hunters. This ~~hunting~~ increase in hunting from the 19th century onwards was an extreme selection pressure that rapidly declined the lynx population because as a predator species, the lynx hasn't evolved to survive purposeful predation by human beings.

The Iberian lynx is also a K-selected species. Females have a low birth rate (only with an average of 3 kits (often decreasing to ≤ 2 after weaning) and parents have to invest significant energy and time into raising the kits. The mother has to catch up to 3 rabbits a day to feed the kits and the kits stay dependent on the parent until 7-10 months and don't leave till 20 months, the young cats also don't begin to mate reproduce until they gain a territory, which can take up to 5 years. These K-type reproductive strategies ~~and~~ the slow sexual maturity and the low litter number (only around 3 a year) of the lynx means that the population grows very slowly and cannot replace itself with the added selection pressure of hunting. Lynx's get killed by hunters at a faster rate than they can reproduce and replace the lost individuals, leading to a decline in the population.

The territoriality of the lynx and the decrease in suitable habitat also contribute to their population decline. The lynx's

hold territories up to 20km^2 and defend from one another. This is to reduce intra-specific competition for space and food, however it also limits the amount of lynx's present in an area. With a huge decline in their heterogeneous habitat, less land is available for lynx's to hold territory. This leads to ~~less~~ displaced individuals ~~not~~ losing their food hunting grounds and starving. Or, individuals are forced to encroach on other lynx's territories, leading to violence between cats and a further reduction in population. Since the lynx's females rarely breed unless they have a territory, less space for territory leads to less females holding territories, leading to less females reproducing and lowering the replacement birthrate of ~~the~~ the population //

The reduction in range and overall population size also makes finding mates more difficult, forcing individuals to mate within their own gene pools and inbreed. This leads to inbreeding depression, ~~where~~ as harmful recessive alleles emerge in inbred individuals, causing detrimental effects to their chances of survival (e.g. no coat growth, poor eyesight). Inbreeding also leads to a reduction in overall genetic diversity. Combined with the effect of genetic drift acting on the ~~by~~ small lynx population, certain alleles can be fixed or lost, reducing the population's fitness in case of sudden ~~or~~ changes in selection pressures. seen

Disease and bacteria are the final factors that have led to the population decline. Due to myxomatosis and haemorrhagic fever epidemics amongst the rabbit populations, the lynx's have lost ~~the~~ large numbers of their main food prey source. A decrease in rabbit leads to more lynx's starving to death and the death of kits as mothers cannot ~~get~~ capture the ≈ 3 rabbits a day needed for the kits energy demands. The outbreak of these diseases also threatens the

health of the lynx population itself. In 2013, it was found that lynx's possess anti-biotic resistant bacteria in their digestive tracts, acquired from the environment. Since the genetic diversity of the lynx population is already low due to inbreeding and the population bottleneck, an outbreak of bacteria and infection could be catastrophic for the species as individuals are less likely to possess alleles to aid their survival during the outbreak due to alleles becoming fixed or lost, threatening extinction for the lynx population. ✓

* ^{seen} The lynx population also underwent a population bottleneck, with only less than 100 animals remaining. This severe reduction in the population causes the bottleneck effect, where alleles from the original gene pool become lost and only a tiny diversity of the alleles remain in the surviving population, increasing their vulnerability to environmental changes and changes in selection pressures. ✓

multiple interventions have been undertaken to avoid the extinction of the species. ✓

The ban protection of the lynx species from hunting in the 1970s would have greatly reduced the deaths of lynx's and relieved them from the species of their main selection pressure. By stopping hunting, less lynx's are killed, slowing the rate of population decline and potentially allowing the birth rate of the species to replace lost individuals and recover its previous size. ✓

The creation of protected areas and the re-introduction of lynx's to previously inhabited areas increases the amount of habitat available to the lynx. By expanding their habitat and spreading individuals out, intraspecific competition amongst lynx's is

reduced among the small isolated populations. Lynx's are given more area to establish adequate territories, which makes more females reproductively viable and increases the birth rate of the species as more females can hold territory. The increased territory also increases the ~~food~~ prey population available to lynx's, meaning that it is more likely that females can catch a sufficient amount of rabbits to feed her kits, reducing the mortality of kits and increasing the population further. 9

The building of highway tunnels and wildlife corridors reduce the likelihood of lynx's being hit by cars at random chance, so they can now cross the increasing amount of highways safely, reducing the amount of individuals killed, preventing further decline. 11

The restocking of rabbits gives the lynx population greater food resources, reducing the deaths by starvation and allowing the population to grow with the additional food sources. 11

The translocation of animals between populations helps to ensure that ~~allele~~ gene flow still occurs and alleles are not lost in isolated populations due to the bottleneck effect and genetic drift, ensuring that the gene pool remains diverse enough to survive changes in the environment. The ~~rest~~ introduction of non-related individuals also reduces the likelihood of inbreeding as new alleles are introduced. This reduces the amount of kits born with genetic defects, improving their chances of survival and leading to more reaching sexual maturity, growing the population.

These interventions have allowed the lynx population to grow from ≈ 100 individuals to 1110 in 2021 ✓

In order to achieve favourable conservation status, the population needs to grow to 3000-3500 individuals. In order to achieve this, the same interventions that were used to grow the population should be continued (the expansion of habitat, re-stocking of rabbits etc), but a few others should be implemented to ensure the long term viability of the population ✓

Populations of rabbits could be treated with ~~anti-~~ anti-biotics, vaccines or the culling of sick individuals to prevent the outbreak of diseases like myxomatosis. Treatment of the rabbit population ensures that the main food source of the lynx is a stable population and that the re-stocked rabbits are not lost to disease. The reduction in these diseases also protect the lynx population from catching ~~diseases~~ ~~to~~ infections from infectal rabbits, further improving the population's chances of survival ✓

Strategies to increase genetic diversity are essential to prevent the effects of a population bottleneck. The 3000-3500 individuals need to have a larger gene pool in order to be less vulnerable to changes in the environment such as disease. Re-introduction of lost alleles could be done through the cloning / unfreezing of preserved lynx dna. Currently in captive breeding programs, lynx embryos have been frozen & for storage. If the genetic diversity of the population is low in the future, genetic alleles stored in these embryos

could be reintroduced into the population to increase allele diversity. since lynx were trophy hunted, preserved tissue of individuals from the original 14th century population might be viable to be used for cloning in order to re-introduce lost alleles back into the lynx population.

Question 2

The moa was the main selection pressure in the environment that caused the evolution of heteroblasty in lancewood due to the moa's herbivory.

moas, ranging in height from that of a turkey to 3m tall, were the main herbivores and largest herbivores in New Zealand's forests. The moa and the lancewood display co-evolution due to the moa's herbivory of the lancewood.

The lancewood displays heteroblasty in order to avoid herbivory by the moa in its early life stages. Grazing from a moa would destroy the plant and potentially take off enough leaves to either kill the plant or severely reduce its growth and chances of survival. Lancewood individuals which were able to avoid herbivory by the moa were able to ~~reduce~~ increase their chances of survival and grow faster without losing energy to ~~growing~~ growing back leaves, increasing

their reproductive success and chance of survival.

The selection pressure of the moa caused directional selection in the lancewood plant. plants that had rigid, barbed phenotypes on their leaves were less likely to be eaten by the moa due to physical harm done to the moa if eaten (cutting of the throat if ingested) compared to soft to eat leaves. Rigid, barbed individuals would have an increased chance of survival, reproducing ~~more~~ more and increasing the frequency of the alleles for stiff, rigid leaves, causing directional selection towards the phenotype seen today. //

Notes

This natural selection would also cause the differences in nutritional content between young and adult leaves. Lancewood plants that ~~produced~~ contained less protein and carbohydrates as a seedling would become ~~more~~ less attractive to herbivores and ~~better~~ conserve their energy by not investing it into their vulnerable leaves. Compared to seedlings with high nutrient density, these plants would be eaten less and lose less energy if a leaf was eaten, increasing their chance of survival and increasing the frequency of those alleles in the gene pool, causing evolution towards heteroblasty. //

The creation and onset of certain random mutations can also help ~~contribute~~ contribute to heteroblasty and increase the lancewood's chances of survival. Random mutations would have occurred, such as the mutation that causes new leaves to be brown in colour, which while either ~~harmful~~ harmful or neutral in other ~~of~~ populations (lack of chlorophyll due to lack of green pigment, reduction in ~~the~~ photosynthesis), greatly increases the lancewood's chances of

Survival due to the camouflage brown leaves ~~proot~~ provide ^{Seen} (*) plants with these ~~colour~~ random mutations (brown leaves and yellow spots on the sides) would be more likely to avoid herbivory by the moa ^m and reach adulthood, increasing the frequency of the mutated allele in the population. In this case, the colouration changes would have become fixed in the population as the allele frequency changed to ~~100%~~ 100%.

After the lancewood reaches a height above ~~that~~ what moas can reach (3 metres for the tallest moa), the plant changes to have the leaves pointed up and begins to resemble a typical tree in the forest. This change in phenotype is due to the selection pressure of the moa disappearing after the plant exceeds a certain height. Without the moa as a selection pressure, the lancewood is subject to other selection pressures such as competition for sunlight and develops a ~~plant~~ phenotype more ~~fit~~ suited to the new ~~set~~ selection pressures (e.g. green leaves to increase photosynthesis). The large difference in selection pressures acting on the lancewood plant between when it's below 3 metres and above 3 metres have caused the plant to develop stark ~~heteroblasty~~ heteroblasty in order to avoid herbivory from the moa species. ^C

^{Seen} (*) Since the brown leaves make the sapling difficult to spot to the moa ^a against a litter of leaves, the moa is unable to spot brown individuals and hence not eat these leaves. Additionally, mutations causing the bright spots on the leaves

edges are highly conspicuous to birds. This may be a form of ~~mimicry~~ mimicry, with the mutation causing the leaves to look poisonous or diseased, deterring birds from eating it //

The lancewood in the mainland and the ~~lance~~ clathrum island lancewood ~~show~~ show ~~a~~ divergent evolution. The two populations are geographically ~~is~~ isolated by sea and have no gene flow between them. Allopatric speciation occurred between the species due to differences in selection pressures between the mainland and clathrum islands. The clathrum islands did not have any browsing birds, so the population there did not have the same selection pressures of herbivory by ~~was~~ moas on the mainland, so the clathrum island lancewood didn't evolve defense mechanisms and heteroblasty to protect itself from predation. The leaves do not change colour and do not have spots or spines or ~~thorns~~ //

~~Reasons~~ One reason the lancewood has retained its heteroblastic lifestyle despite the extinction of the moa is due to the moa not being extinct for long enough. The moa only went extinct around 600 years ago. The timeframe of large-scale evolutionary changes is much longer than 600 years. Within 600 years, there hasn't been enough generations of the species for the previously unfavourable alleles (such as green sapling colouration and soft leaves) that would previously been selected against to re-emerge in the population and increase in allele frequency to cause meaningful changes to the species. //

Additionally, there aren't any strong selection pressures against the heteroblastic life cycle of lancewood. Traits such as downwards facing leaves, bright spots and tough spines are not directly harmful to the survival of the lancewood. Without selection pressures against these alleles, the allele frequency of the population will remain the same with the exception of genetic drift, which is random and ~~not fast~~ doesn't occur fast enough (especially in a large population) to have any noticeable ~~effect~~ effect on the evolution of the lancewood within 600 years. //

Certain alleles such as those for green ~~sep~~ coloured saplings and soft leaves might have been lost in the gene pool due to the selection pressure of the moa, with ~~some~~ individuals carrying those alleles being eaten by moa and outcompeted by ~~better~~ better adapted plants, those alleles could be lost, meaning that the only way for new alleles to be introduced into the lancewood ~~to~~ population is through random mutation, which is unpredictable ~~and~~ and slow. //

The last reason the lancewood ~~could~~ is retaining retaining its heteroblastic life cycle ~~to~~ could be the introduction of new browsing herbivores into the NZ ecosystem. After moas went extinct, ~~deer~~ deer, goats and pigs were introduced to New Zealand. Wild populations of these

herbivores could have replaced the ~~ecological~~ ecological niche of ~~the~~ the now once inhabited. ~~As~~ since deer are large, browsing herbivores, they continue to exert selection pressures on the lancewood population to retain ~~its~~ ^{their} defensive mechanisms as saplings. Any lancewood plants without heteroblasty or with unfavourable traits like high nutrient dense leaves or no-spots might be instead eaten by deer and other mammals, preventing those alleles from being re-introduced back into the lancewood population. //

Question 34

the biological species concept only works effectively on sexually reproducing populations where distinct and obvious reproductive isolation is present between populations. Many times species are described off geographical separation and differences in phenotype, however when species are brought together, the reproductive isolation ~~is~~ is revealed to not exist and the populations hybridize ~~and~~ with each other.

The main example of this is between the parula and mallard duck species. Geographically they were very isolated, with the mallard originating from open grassland in North America the northern hemisphere while the parula is endemic to NZ (although closely related to other duck species in the Pacific). ~~Extremely~~ externally, their phenotypes are also clearly different, with the mallard having a distinctly differently green head in males. However, their reproductive behavior is still very similar. Both species have overlapping breeding and egg laying months, and clearly parula and mallards do recognize each other as mates. This means that there are no little pre-zygotic reproductive isolating mechanisms and evidently, neither do post-zygotic reproductive isolating mechanisms as hybrids are fertile and do not experience hybrid breakdown. As soon as the species are inhabiting the same range (NZ), previously assumed RIMs ~~cause~~ cease to exist as clearly the populations haven't been

Isolated geographically for long enough to develop other RIMs, such as behavioural RIMs or post-zygotic RIMs (different chromosome number or gamete incompatibility incompatibility). ✓

Similar examples are seen ~~as~~ elsewhere. In North America, grizzly bears and polar bears ~~remained~~ ^{remained} separate populations, with the polar bears ~~solely~~ ^{solely} inhabiting the arctic ice. However with climate change, the polar bears are losing habitat and are forced further south to escape melting ice sheets, coming into contact with grizzly bears. More and more hybrids between grizzly and polar bears are being sighted with both white and brown fur colourations.

Orcas and shortfinned pilot whales also often create hybrids ~~due~~ due to the two species, while ~~so~~ starkly different in colouration, socialising together and cooperatively hunting. Although the species seem different, there are basically no behavioural RIMs ~~separating~~ separating the ~~two~~ two ~~as~~ blackfish species.

The concept of a species is especially ~~very~~ blurry when applied to tropical reef fishes, especially those like the pipefish family. Fish from groups often rapidly speciate due to limited range ~~over~~ reef (separate reefs) and change colouration dramatically. However when species are put in the same ~~habitat~~ ^{habitat}, hybridisation ~~occurs~~ ^{occurs} freely and ~~many~~ ^{many} species are able to change their phenotype (colours) at will to better ~~suit~~ ^{suit} their new environment, making fish reef fish species incredibly hard to differentiate from the naked eye of a human. All these examples

show how very ~~a~~ obvious differences to our perception, like range and phenotype, are only surface level and certain special groups/species can overcome RIMS that totally isolate other populations from each other.

Alternative evidence to distinguish the mallard and purebred species aside from ~~pure~~ external soft-body features (feathers, colours, etc) include genetic markers, internal or hard-body features, and differences in behaviour.

Definitive genetic studies have yet to be undertaken with either species. During their geographic separation, it is very likely that mutations occurred in the genomes of either species that are not present in the other. While these mutations might have no effect on RIMS, they could be used to determine a duck's genetic purity. —
 mtDNA analysis could be used to examine point mutations on the mtDNA of purebred to determine which individuals are descended from the original purebred population. However, mtDNA is passed solely from the mother so a hybrid could possess 100% purebred mtDNA if the female was purebred. Nuclear DNA could be analysed to find unique mutations and alleles present in the purebred population. Heterozygous individuals for these alleles would most likely be pure-bred purebred as only purebred parents could pass those alleles. These genetic markers, while unrecognizable to the average person, could help conservationists locate and isolate purebred purebred individuals. 4

The ~~its~~ soft-part and plumage features of pure and hybrids are hard to distinguish and harder to when it's a hybrid. However, less obvious features like skeletal structure could be used to ~~determine~~ distinguish species, as these features are hidden from plain view and could reveal more obvious differences/variation between the two species. ✓


lastly, certain behaviours or vocalizations of either species could be analyzed for differences. A pure-mallard hybrid might use vocalizations that are an obvious mix of the two species due to competing by both a mallard and a pure parent. Ducks that only had pure parents might have a distinct vocalization that can be verified for authenticity. ✓


The difficulties with the species concept between mallards and pure impacts decisions on conservation efforts for the pure. ✓


Firstly, due to the rampant hybridization and difficulty distinguishing between the 2 species, the shooting of pure by game hunters cannot be reliably policed and prevented. since the mallard is very common, hunters are free to shoot them & however the critically endangered pure can be thrown into that mix and individuals could be lost without the overarching protection from conservation organizations. Additionally, the genetic mixing


between the ducks creates tricky legal and biological implications if the shooting of ~~pure~~ pure parera becomes regulated/banned, because at what point is a hybrid considered pure? If the hybridization is widespread, there might be no individual ducks that can be accurately classified as ~~endangered~~ "endangered" or "free to shoot", making the regulation of hunting infeasible. 9


As secondly, with very few/no pure parera left, efforts to maintain a genetically intact population for conservation are further complicated. Establishing only pure breeding populations or only might lead to the species detriment with so few individuals, inbreeding depression and ~~population~~ the bottleneck effect might decrease the fitness of the already critical pure population. The declining number of pure also pose a ~~an~~ unclear question to conservationist. Perhaps it is ~~now~~ better to preserve the genetic history of the pure species by allowing it to hybridize with a more successful species and keep the genes in the now mixed population rather than maintain a pure breeding population that is vulnerable to extinction, losing the pure genetic history forever. In the future, the now hybridized population of pure might become the only remains of the original pure species, which may cause science to now


Classify the hybrids as the new purebred population
and these nullard-purebred hybrids receive the
full ~~pro~~ protection of a native species in
NR. perhaps the protection would have to extend
to the entire nullard population 














93101A

