



S-CALCF



Mana Tohu Mātauranga o Aotearoa
New Zealand Qualifications Authority

Scholarship 2024

Calculus

FORMULAE AND TABLES BOOKLET

Refer to this booklet to answer the questions for Scholarship Calculus 93202.

Check that this booklet has pages 2–4 in the correct order and that none of these pages is blank.

YOU MAY KEEP THIS BOOKLET AT THE END OF THE EXAMINATION.

CALCULUS – USEFUL FORMULAE

ALGEBRA

Quadratics

If $ax^2 + bx + c = 0$

then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Logarithms

$y = \log_b x \Leftrightarrow x = b^y$

$\log_b(xy) = \log_b x + \log_b y$

$\log_b\left(\frac{x}{y}\right) = \log_b x - \log_b y$

$\log_b(x^n) = n \log_b x$

$\log_b x = \frac{\log_a x}{\log_a b}$

Binomial Theorem

$$(a+b)^n = \binom{n}{0}a^n + \binom{n}{1}a^{n-1}b^1 + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + \binom{n}{n}b^n$$

$$\binom{n}{r} = {}^nC_r = \frac{n!}{(n-r)!r!}$$

Some values of $\binom{n}{r}$ are given in the table below.

| $\begin{smallmatrix} n \\ \backslash \\ r \end{smallmatrix}$ | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|--|---|----|----|-----|-----|-----|-----|-----|-----|-----|----|
| 0 | 1 | | | | | | | | | | |
| 1 | 1 | 1 | | | | | | | | | |
| 2 | 1 | 2 | 1 | | | | | | | | |
| 3 | 1 | 3 | 3 | 1 | | | | | | | |
| 4 | 1 | 4 | 6 | 4 | 1 | | | | | | |
| 5 | 1 | 5 | 10 | 10 | 5 | 1 | | | | | |
| 6 | 1 | 6 | 15 | 20 | 15 | 6 | 1 | | | | |
| 7 | 1 | 7 | 21 | 35 | 35 | 21 | 7 | 1 | | | |
| 8 | 1 | 8 | 28 | 56 | 70 | 56 | 28 | 8 | 1 | | |
| 9 | 1 | 9 | 36 | 84 | 126 | 126 | 84 | 36 | 9 | 1 | |
| 10 | 1 | 10 | 45 | 120 | 210 | 252 | 210 | 120 | 45 | 10 | 1 |
| 11 | 1 | 11 | 55 | 165 | 330 | 462 | 462 | 330 | 165 | 55 | 11 |
| 12 | 1 | 12 | 66 | 220 | 495 | 792 | 924 | 792 | 495 | 220 | 66 |

Complex numbers

$z = x + iy$

$= r \operatorname{cis} \theta$

$= r(\cos \theta + i \sin \theta)$

$\bar{z} = x - iy$

$= r \operatorname{cis}(-\theta)$

$= r(\cos \theta - i \sin \theta)$

$r = |z| = \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}$

$\theta = \arg z$

where $\cos \theta = \frac{x}{r}$

and $\sin \theta = \frac{y}{r}$

De Moivre's Theorem

If n is any integer, then

$(r \operatorname{cis} \theta)^n = r^n \operatorname{cis}(n\theta)$

COORDINATE GEOMETRY

Straight Line

Equation $y - y_1 = m(x - x_1)$

Circle

$(x - a)^2 + (y - b)^2 = r^2$

has a centre (a, b) and radius r

Parabola

$y^2 = 4ax$ or $x = at^2$ $y = 2at$

Focus $(a, 0)$ Directrix $x = -a$

Ellipse

$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ or $x = a \cos \theta$ $y = b \sin \theta$

Foci $(c, 0), (-c, 0)$ where $b^2 = a^2 - c^2$

Eccentricity: $e = \frac{c}{a}$

Hyperbola

$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ or $x = a \sec \theta$ $y = b \tan \theta$

asymptotes $y = \pm \frac{b}{a}x$

Foci $(c, 0), (-c, 0)$ where $b^2 = c^2 - a^2$

Eccentricity: $e = \frac{c}{a}$

CALCULUS

Differentiation

| $y = f(x)$ | $\frac{dy}{dx} = f'(x)$ |
|--------------------------|----------------------------------|
| $\ln x$ | $\frac{1}{x}$ |
| e^{ax} | ae^{ax} |
| $\sin x$ | $\cos x$ |
| $\cos x$ | $-\sin x$ |
| $\tan x$ | $\sec^2 x$ |
| $\sec x$ | $\sec x \tan x$ |
| $\operatorname{cosec} x$ | $-\operatorname{cosec} x \cot x$ |
| $\cot x$ | $-\operatorname{cosec}^2 x$ |

Integration

| $f(x)$ | $\int f(x)dx$ |
|----------------------|--|
| x^n | $\frac{x^{n+1}}{n+1} + c$ ($n \neq -1$) |
| $\frac{1}{x}$ | $\ln x + c$ |
| $\frac{f'(x)}{f(x)}$ | $\ln f(x) + c$ |

First principles

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Parametric Function

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$$

$$\frac{d^2y}{dx^2} = \frac{d}{dt} \left(\frac{dy}{dx} \right) \cdot \frac{dt}{dx}$$

Product Rule

$$(f \cdot g)' = g \cdot f' + f \cdot g' \text{ or if } y = uv \text{ then } \frac{dy}{dx} = v \frac{du}{dx} + u \frac{dv}{dx}$$

Quotient Rule

$$\left(\frac{f}{g} \right)' = \frac{g \cdot f' - f \cdot g'}{g^2} \text{ or if } y = \frac{u}{v} \text{ then } \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

Composite Function or Chain Rule

$$(f(g))' = f'(g) \cdot g'$$

$$\text{or if } y = f(u) \text{ and } u = g(x) \text{ then } \frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

NUMERICAL METHODS

Trapezium Rule

$$\int_a^b f(x) dx \approx \frac{1}{2} h [y_0 + y_n + 2(y_1 + y_2 + \dots + y_{n-1})]$$

$$\text{where } h = \frac{b-a}{n} \text{ and } y_r = f(x_r)$$

Simpson's Rule

$$\int_a^b f(x) dx \approx \frac{1}{3} h [y_0 + y_n + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$$

$$\text{where } h = \frac{b-a}{n}, y_r = f(x_r) \text{ and } n \text{ is even.}$$


TRIGONOMETRY

$$\operatorname{cosec} \theta = \frac{1}{\sin \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\cot \theta = \frac{1}{\tan \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}$$

**Sine Rule**

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Cosine Rule

$$c^2 = a^2 + b^2 - 2ab \cos C$$

Identities

$$\cos^2 \theta + \sin^2 \theta = 1$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$\cot^2 \theta + 1 = \operatorname{cosec}^2 \theta$$

General Solutions

$$\text{If } \sin \theta = \sin \alpha \text{ then } \theta = n\pi + (-1)^n \alpha$$

$$\text{If } \cos \theta = \cos \alpha \text{ then } \theta = 2n\pi \pm \alpha$$

$$\text{If } \tan \theta = \tan \alpha \text{ then } \theta = n\pi + \alpha$$

where n is any integer

Compound Angles

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

Double Angles

$$\sin 2A = 2 \sin A \cos A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

$$\begin{aligned} \cos 2A &= \cos^2 A - \sin^2 A \\ &= 2 \cos^2 A - 1 \\ &= 1 - 2 \sin^2 A \end{aligned}$$

Products

$$2 \sin A \cos B = \sin(A+B) + \sin(A-B)$$

$$2 \cos A \sin B = \sin(A+B) - \sin(A-B)$$

$$2 \cos A \cos B = \cos(A+B) + \cos(A-B)$$

$$2 \sin A \sin B = \cos(A-B) - \cos(A+B)$$

Sums

$$\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$$

$$\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$$

$$\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}$$

$$\cos C - \cos D = -2 \sin \frac{C+D}{2} \sin \frac{C-D}{2}$$

MEASUREMENT**Triangle**

$$\text{Area} = \frac{1}{2} ab \sin C$$

Trapezium

$$\text{Area} = \frac{1}{2} (a+b)h$$

Sector

$$\text{Area} = \frac{1}{2} r^2 \theta$$

$$\text{Arc length} = r\theta$$

Cylinder

$$\text{Volume} = \pi r^2 h$$

$$\text{Curved surface area} = 2\pi rh$$

Cone

$$\text{Volume} = \frac{1}{3} \pi r^2 h$$

$$\text{Curved surface area} = \pi rl \text{ where } l = \text{slant height}$$

Sphere

$$\text{Volume} = \frac{4}{3} \pi r^3$$

$$\text{Surface area} = 4\pi r^2$$